GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: **B6J.129S6-*Sptlc1*$^{tm1.1Rlp}$/Mmucd** **MMRRC:** **042294-UCD**

Protocol:

GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM)	0.5
Primer 2. (stock concentration is 20µM)	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	**15.00 µL**

Comments on protocol:

- Protocol may work with other DNA extraction methods
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65ºC decreasing in temperature by 1.0ºC; next 30 cycles anneal at 55ºC.

Strategy:

Steps		Temp (ºC)	Time (m:ss)	# of Cycles
1. Initiation/Melting	HOT START? ☐	94	5:00	**1**
2. Denaturation		94	0:15	
3. Annealing	steps 2-3-4 cycle in sequence	65 to 55 (↓1ºC/cycle)	0:30	**40x**
4. Elongation		72	0:40	
5. Amplification		72	5:00	**1**
6. Finish		15	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1.42294-loxF	GGGTTCTATGGCACATTTGGTAAG
2.42294-loxR	CTGTTACTTCTTGCCAGTGGAC

Electrophoresis Protocol:

Agarose: **1.5%** V: **90**

Estimated Running Time: **90** min.

Primer Combination	**Band** (bp)	Genotype
1 & 2	350	wildtype
	425	LOX



PCR protocol developed by MMRRC at University of California, Davis