

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR1431 2810474O19Rik EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	3.9
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.6
Primer 2. (stock concentration is 20µM) wtR	1.2
Primer 3. (stock concentration is 20µM) mutR	0.3
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_2810474O19Rik-comF	GTAAGCTTTGCTTTACAAC TAGCATGG	Estimated Running Time: 90 min.	
2. CR_2810474O19Rik-wtR	GATGACTGGGTTTTAGAATATGGTGG	Primer Combination	Band (bp)
3. CR_2810474O19Rik-mutR	CTGCCCTATTTACTCAGCCACTCC	1 & 2, 1 & 3	610,5087*
		1 & 3	238
			Genotype
			wildtype
			mutant

Electrophoresis Protocol:

Allele Description: Exon 3 [ENSMUSE00000315731](#) and flanking splicing regions were constitutively deleted from the 2810474O19Rik gene [ENSMUST0000046689.12](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

*May not see larger wildtype band

