

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

Protocol Name: CR1635 Prr9 GenDel

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
<b>TOTAL VOLUME OF REACTION:</b>	<b>15.00 µL</b>

### Comments on protocol:

- Protocol may work with other DNA extraction methods.

### Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting <span style="float: right;">HOT START? <input type="checkbox"/></span>	94	2:00	<b>1x</b>
2. Denaturation	94	0:10	
3. Annealing <span style="float: right;">steps 2-3-4 cycle in sequence</span>	65 (↓1°C/cycle)	0:30	<b>10x</b>
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing <span style="float: right;">steps 5-6-7 cycle in sequence</span>	55	0:30	<b>25x</b>
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

### Primers:

Name	Nucleotide Sequence (5' - 3')	Electrophoresis Protocol:		
1. CR_Prr9-comF	GGCTGATTAACAGGCATCACAGC	Agarose: 1.5%	V: 90	
2. CR_Prr9-wtR*	TGTCCAGTCCTCCACAAAATCTAAT	Estimated Running Time: 90 min.		
3. CR_Prr9-mutR	ACCATTTTTGATTTTGTAGCATCACC	<b>Primer Combination</b>	<b>Band (bp)</b>	<b>Genotype</b>
		1 & 2, 1 & 3	545, 1583	wildtype
		1 & 3	326	mutant

**Allele Description:** Exon 1-2 ([ENSMUSE00000420276](#), [ENSMUSE00000420269](#)) were constitutively deleted between the 5'UTR through the 3' UTR from the Prr9 gene [ENSMUST0000070284.3](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

\*wtR primer untested (ePCR verified)

