

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

Protocol Name: CR1568 Ms4a12 iDex

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
<b>TOTAL VOLUME OF REACTION:</b>	<b>15.00 µL</b>

### Comments on protocol:

- Protocol may work with other DNA extraction methods.

### Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting <span style="float: right;">HOT START? <input type="checkbox"/></span>	94	2:00	<b>1x</b>
2. Denaturation	94	0:10	
3. Annealing <span style="float: right;">steps 2-3-4 cycle in sequence</span>	65 (↓1°C/cycle)	0:30	<b>10x</b>
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing <span style="float: right;">steps 5-6-7 cycle in sequence</span>	55	0:30	<b>25x</b>
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

### Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Ms4a12-comF	GTATGTGCTCACTTGCTCTCTCTCC	Estimated Running Time: 90 min.	
2. CR_Ms4a12-wtR*	CTGTGTGTTAGGGAAGATTCCTGG	<b>Primer Combination</b>	<b>Band (bp)</b>
3. CR_Ms4a12-mutR	ATGTGTTTGCTCATTGGTTGATATGC	1 & 2, 1 & 3	295,476
		1 & 3	295
			wildtype
			mutant

### Electrophoresis Protocol:

**Allele Description:** Exon 1 [ENSMUSE00001330093](#) had 181bp deleted from the 33<sup>rd</sup> coding nucleotide through the 213<sup>th</sup> coding nucleotide from the Ms4a12 gene [ENSMUST00000186228.1](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

\*Run in Simplex

\*wtR primer untested (ePCR verified)

