

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

Protocol Name: CR10057 Fkbp11 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
<b>TOTAL VOLUME OF REACTION:</b>	<b>15.00 µL</b>

**Comments on protocol:**

- Protocol may work with other DNA extraction methods.

**Strategy:**

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting <span style="float: right;">HOT START? <input type="checkbox"/></span>	94	2:00	<b>1x</b>
2. Denaturation	94	0:10	
3. Annealing <span style="float: right;">steps 2-3-4 cycle in sequence</span>	65 (↓1°C/cycle)	0:30	<b>10x</b>
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing <span style="float: right;">steps 5-6-7 cycle in sequence</span>	55	0:30	<b>25x</b>
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

**Primers:**

**Electrophoresis Protocol:**

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR-Fkbp11_comF	AGACACGCTCCACATACACTACACG	Estimated Running Time: 90 min.	
2. CR-Fkbp11_wtR*	CCATCTACCAAAGTCCCTGAAGAC	<b>Primer Combination</b>	<b>Band (bp)</b>
3. CR-Fkbp11_mutR	CTGCACCTCTCCAGTATGAAGTCAGG	1 & 2, 1 & 3	426,934
		1 & 3	525
			wildtype
			mutant

**Allele Description:** Exon 3 [ENSMUSE00000132540](#) and flanking splicing regions were constitutively deleted from the Fkbp11 gene [ENSMUSG0000003355](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

\*wtR primer untested (ePCR verified)

