

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR10201 Lsm8 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Lsm8_comF	TGAAGGAAAGTAGAGGTGATTCATGGAG	Estimated Running Time: 90 min.	
2. CR_Lsm8_wtR*	CTGTAGTCCAGAAGTATTCCACGACACA	Primer Combination	Band (bp)
3. CR_Lsm8_mutR	TGCATCGTGGCACACATTAAGTACAT	1 & 2, 1 & 3	336, 1188
		1 & 3	373
			wildtype
			mutant

Allele Description: Exon 3 [ENSMUSE00001297276](#) and flanking splicing regions were constitutively deleted from the Lsm8 gene [ENSMUSG00000044155](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

*wtR primer untested (ePCR verified)

